

# Effect of Butternut Squash (*Cucurbita Moschata*) Seeds Powder on the Chemical and Rheological Properties of Stirred Cultured Camel Milk and Yoghurt

Collins K. Kangogo, Patrick S. Muliro and Joseph O. Anyango\*

Department of Dairy, Food Science and Technology, Egerton University, Nakuru, Kenya.

\* Corresponding author: ajochieng@egerton.ac.ke

## Abstract

Stirred fermented camel milk products are difficult to produce because of their weak texture and thin consistency, which makes them unpopular with consumers. Research conducted on camel milk revealed that during the coagulation process, only flocs were formed without firm coagulation, attributed to the lack of interaction between  $\kappa$ -casein and  $\beta$ -lactoglobulin, the large casein micelle size, leading to a weak network of casein formation. In an effort to address this issue, researchers turned to corn starch as a thickening agent, discovering that a concentration of 2.0% effectively improved the viscosity and significantly reduced syneresis in stirred camel milk yoghurt and stirred cultured camel milk. However, this study seeks to explore alternatives to corn starch, given its predominant role as a staple food in Kenya and its multifaceted applications across various industries. Butternut squash seeds being a by-product, presents a promising substitute to corn starch, given its abundant hydrocolloid composition. By incorporating butternut squash (*Cucurbita moschata*) seed powder (BSSP) as a thickener, this study aims to enhance the chemical and rheological properties of stirred camel milk yoghurt and stirred cultured camel milk. Stirred fermented camel milk product were prepared using 4 litres of camel milk, 2% starter cultures (thermophilic culture for yoghurt and mesophilic aromatic culture for stirred cultured camel milk) and BSSP 0.0% (negative control), 0.4%, 0.8%, 1.2%, 1.6%, 2.0% mixed with 0.4% gelatin. 2.0% corn starch mixed with 0.4% gelatin was used as a standard for comparison. Results show that increasing the BSSP level significantly ( $p < 0.05$ ) decreased the moisture content while increasing the total solid content of stirred fermented camel milk products. There was a progressive increase in ash content with increase in BSSP levels. There was a significant ( $p < 0.05$ ) reduction in the pH, with increase in BSSP levels in stirred fermented camel milk samples. Increasing the concentration of BSSP from 0.4% to 2.0% resulted in significant ( $p < 0.05$ ) increase in viscosity and a reduction in syneresis of stirred camel milk yoghurt and stirred cultured camel milk samples. This study demonstrates that BSSP effectively enhances the viscosity, reduces syneresis and increases acidity in stirred fermented camel milk products during storage.

**Keywords:** Corn starch, Butternut squash seed powder, Gelatin, stirred camel milk yoghurt, Stirred cultured camel milk.

## 1. INTRODUCTION

The limited availability of processed camel milk products in Kenya can be attributed to technical challenges and low consumer receptiveness. This is because it is challenging to replicate the viscosity of camel milk to match the familiar consistency of bovine milk that customers are accustomed to. Stirred fermented camel milk products are difficult to produce because camel milk does not coagulate well, resulting in a thin texture and frail product structure, with the gel texture being an important property in determining the mouthfeel, appearance and overall acceptability by

consumers in the modern market with a shift in preference to a particular texture, such as thicker mouthfeel (Seifu, 2023; Wong *et al.*, 2020). The weak firmness of camel milk coagulum can be attributed to lack of  $\kappa$ -casein and  $\beta$ -lactoglobulin interactions (Ho *et al.*, 2022); the large casein micelle size and low amounts of  $\kappa$ -casein and high ratio of whey proteins to caseins, resulting in a fragile casein network that can easily be disrupted during coagulum formation (Berhe *et al.*, 2017).

Various dairy constituents (for example, whey protein concentrate) and hydrocolloids (for example starch and various gums) have been utilized to combat defects in fermented milk like syneresis, poor body and texture (Hematyar *et al.*, 2012). Hydrocolloids are non-digestible carbohydrates that have the ability to form gels or viscous solutions, thickening, emulsifying, coating and stabilizing (Hafezi & Pirsá 2023). Hydrocolloids consist of lengthy and complex molecules that can either interconnect internally or with other molecules in their surroundings to create an emulsion. They are hydrophilic polymers that have many hydroxyl groups and may be polyelectrolytes (Abdelmoneim *et al.*, 2016). These polysaccharides work by interacting with the casein network and contribute to the formation of gels in a two-phase system (liquid-solid); a continuous, solid, three dimensional network structure forming the gel matrix holding finely dispersed liquid phase (Lunardello *et al.*, 2012). Addition of hydrocolloids maintains the morphological characteristics of yoghurt during transportation and storage by increasing its viscosity, absorbing water, strengthening and improving its texture. They are derived from animals, plants, microorganisms, or synthetic origins and are naturally present in foodstuffs (Pirsá & Hafezi, 2023). There is a rising need for hydrocolloids because of their capacity to create a stable emulsion system in dairy products, preventing whey separation and protein clumping throughout their shelf life (Javidi *et al.*, 2016). Through various mechanisms like covalent, electrostatic, excluded volume, hydrophobic interactions, van der waal forces, hydrocolloids combine with milk proteins to establish a gel network system, thereby enhancing the texture of dairy items (De Kruijff and Tuinier, 2001; Wusigale *et al.*, 2020).

Only a small portion of Butternut squash is utilized directly for human consumption whereas the remaining portion (skin and seeds) is undervalued and treated as by-products and waste. These wastes can make important and valuable contribution to food resources and development of industrial products considering the abundant nutrition and technological properties contained in these residues (Li, 2020). A major issue encountered by many nations is handling waste especially from food products and agricultural produce. These waste has significant global environmental impact due its biodegradable nature and high water content contributing to pollution and has the potential to produce bad odour, abundant leachate (Liu *et al.*, 2012) and emit greenhouse gases (for example methane) in landfills (Cheng & Hu, 2010).

Butternut seeds boast a rich nutrition profile, featuring lipids, proteins, vitamins, phytosterols, minerals such as potassium, calcium and sodium and trace elements (Jiao *et al.*, 2014; Sztark *et al.*, 2010). Specifically, butternut squash seeds have a protein content of 32.38%, a fat content of 14.31% and a total phenolic content of  $1.82 \pm 0.05$  mg GAE/g (Li, 2020). It also contains a fibre content  $1.152 \pm 0.20\%$  and ash content of  $1.152 \pm 0.20\%$  (Assilidjoe *et al.*, 2021). Wang *et al.* (2017) obtained polysaccharides from defatted butternut squash seed powder, the yield of crude polysaccharide was 2.3 %. Hydrocolloids from Butternut squash (*Cucurbita moschata*) seeds were evaluated as a stabilizer to preserve the physicochemical properties, flavour and texture of yoghurt samples. The addition of hydrocolloids from butternut squash seeds modified the rheology with positive effects on the texture, improved the physical stability and overall mouthfeel properties to

increase desirable sensorial attributes without any consequent appearance of undesirable effects (Rojas-Torres *et al.*, 2021). The researchers concluded that hydrocolloids from butternut squash seeds could be used as an additive for the formulation of microstructure food matrices. It can be utilized alone or blended with other hydrocolloids to increase viscosity or thickness and improve the water binding ability and texture of yoghurt.

There have been several attempts to remedy these challenges facing production of stirred fermented camel milk products. The reported approaches include using different gelling and thickening agents either alone or together (for example gelatin, corn starch, modified starch, bovine skim milk powder, polymerized whey protein isolate), hydrocolloids (for example xanthan gum, pectin, gum Arabic, alginate, gum acacia carboxymethyl cellulose,  $\kappa$ -carrageenan, sodium carboxymethyl cellulose) and other stabilizers such as calcium chloride  $\text{Na}_2\text{EDTA}$  and mono- and di-glyceride (Al-Otaibi & Al-Zoreky, 2015; Galeboe *et al.*, 2018; Hashim *et al.*, 2009; Ho *et al.*, 2022; Ibrahim & Khalifa; Jasim *et al.*, 2018; Kavas, 2016; 2015; Muhammad Imran, 2014; Muliro *et al.*, 2013; Oselu *et al.*, 2022).

This study examined how varying concentrations (0.4, 0.8, 1.2, 1.6, 2.0%) of BSSP affected the chemical composition of both stirred stirred camel milk yoghurt and stirred stirred cultured camel milk. The rheological characteristics of stirred camel milk yoghurt and stirred cultured camel milk was also studied for the 21 days of storage at 1, 7, 14, 21 days interval. Two controls were employed for this experiment; the negative control consisting of camel milk mixed with 0.4% gelatin and a standard consisting of camel milk mixed with 2.0% corn starch and 0.4% gelatin.

## **2. MATERIALS AND METHODS**

### **2.1 Sample Collection**

Fresh camel milk was procured from Marigat, Baringo county, Kenya. Fresh raw camel milk was conveyed in clean 20 litre containers to Department of Dairy and Food Science and Technology, Egerton University, Nakuru City, Kenya. Gelatin powder (beef skin), corn starch (Tri-Clover industries Kenya Limited) were all procured from Gilanis supermarket, Nakuru city, Kenya. Starter cultures (thermophilic and mesophilic) were procured from Promaco Kenya limited. Whole butternut squash seeds were obtained Vegpro Kenya Limited, Nairobi city, Kenya.

### **2.2 Preparation of Butternut Squash (*Cucurbita moschata*) Seeds**

Fresh butternut squash seeds were collected from Vegpro Kenya limited during butternut squash processing, cleaned with distilled water and then oven dried at 65 °C for 12 hours as described by Li, (2020). The seeds were then powdered using a hammer mill and subsequently sieved using a no.30 sieve (600 $\mu\text{m}$ ) in size to remove any coarse particles. Powdered butternut squash seeds were stored in a zip lock bags until use.

### **2.3 Preparation of Stirred fermented camel milk Products**

Prior to making of stirred fermented camel milk products, all equipment was carefully cleaned and sterilized using hot water. Different formulations were employed to produce stirred fermented camel milk products as indicated in **Table 1**; this was achieved by altering the concentrations of

BSSP (0.4, 0.8, 1.2, 1.6, 2.0%), while maintaining the concentration of gelatin at 0.4%. Fresh camel milk mixed with 0.4% gelatin was utilized as the negative control and fresh camel milk mixed with 2.0% corn starch and 0.4% gelatin as the standard. Stirred fermented camel milk products (yoghurt and stirred cultured camel milk) was produced using steps outlined by Muliro *et al.*(2013); Oselu *et al.* (2022) but with minor changes. For stirred camel milk yoghurt samples, raw camel milk was pre heated to 65°C to facilitate addition of dry ingredients (stabilizers). The mixture was then pasteurized in a thermostatically controlled water bath at 85°C for 30 minutes then allowed to cool to 45°C to facilitate inoculation of thermophilic yoghurt culture (Chr Hansen, YF-L903, Denmark) which consisted of strains of *Lactobacillus delbrueckii subsp. bulgaricus* and *Streptococcus thermophilus*. Incubation was then carried out at 45°C for 6h and refrigerated at 4°C until the samples were analyzed. For stirred cultured camel milk samples, raw camel milk was pre heated to 65°C to facilitate addition of dry ingredients (stabilizers). The mixture was then pasteurized at 85°C for 30 minutes. The pasteurized mixture was then cooled to 25°C to facilitate inoculation of mesophilic aromatic culture (Chr Hansen, CHN-22, Denmark) which consisted strains of *Lactococcus lactis subsp. Cremoris*, *Leuconostoc*, *Lactococcus lactis subs. lactis*, *Lactococcus lactis subs. lactis biovar diacetylactis*. Fermentation was carried out at 25°C for 24h.

## 2.3 Chemical Properties

### 2.3.1 Determination of Total Solid Content

The weight of the residue obtained from analysis of moisture content in stirred fermented camel milk samples was used to calculate the total solids using the formula in equation 1 as described by Association of Official Analytical Chemists (AOAC, 2005, Method 925.23):

$$\text{Total Solids (\%)} = \frac{\text{dry sample}}{\text{weight of the sample}} \times 100 \dots \text{Equation 1}$$

### 2.3.2 Determination of Ash Content

The ash content of the stirred fermented camel milk products was determined following the AOAC (2005 Method number 930.05). 10 g of stirred fermented camel milk samples were weighed and transferred to a dry crucible. The samples were subjected to charring using a flame to decompose all organic components. Subsequently, the samples were ashed in a muffle furnace and incinerated at 550 °C for 6 h and cooled in a desiccator for 10 mins, the samples were weighed, and the ash content was calculated using the formula in equation 2:

$$\text{Ash(\%)} = \frac{\text{weight of residue}}{\text{weight of sample}} \times 100 \dots \text{Equation 2}$$

### 2.3.3 Determination of pH

The pH of stirred fermented camel milk products was determined using a pH meter (Adwa Instruments, Model 1020, Szeged, Hungary). The meter was calibrated using standard buffer solution of pH 4.0 and pH 7.0 and rinsed with distilled water.

## 2.4 Rheological Properties

### 2.4.1 Measurement of Syneresis

10ml of gelled stirred fermented camel milk product was centrifuged at 3000 rpm for 10 minutes at 10°C according to (Muliro *et al.*, 2013) with slight modifications. The syneresis of stirred fermented camel milk products was determined at 1, 7, 14, 21 days. Measurement of susceptibility to syneresis was determined using the formula in equation 3:

$$\text{Syneresis}(\%) = \frac{\text{Volume of separated whey}}{10\text{ml of fermented camel milk product}} \times 100\% \dots\dots\dots \text{Equation 3}$$

### 2.4.2 Measurement of Viscosity.

The viscosity of stirred fermented camel milk products was determined at 10°C for stirred camel milk yoghurt and 20°C for stirred cultured camel milk using DVELV viscometer (Brookfield Ametek, Model LV, Middlesboro, USA). Spindle no.63 and 20rpm (rotation speed) was used to measure 500 ml of stirred fermented camel milk products. Three replicates of stirred camel milk yoghurt and stirred cultured milk treatments were taken in centipoise units (cP)

### 2.5 Data Analysis

Data analysis was carried out using Statistical Analysis System (SAS Institute Inc., 2006) software Version 9.4 and Microsoft Excel (Version 2016). To test the study hypothesis, analysis of variance (ANOVA) was performed. The level of significance was established at P<0.05 confidence level and means separation was done using Tukey’s Honest significant difference (HSD) method.

## 3. RESULTS AND DISCUSSION

### 3.1 Effect of BSSP on the Chemical Properties of Stirred Fermented Camel Milk Products

#### 3.1.1 Total Solid Content

The total solid content values for stirred camel milk yoghurt samples ranged from 10.90% to 15.78% as shown in Table 1. Whereas stirred cultured camel milk samples ranged from 11.03% to 14.67% as shown in Table 2. Addition of BSSP significantly (p<0.05) increased the total solids content of stirred camel milk yoghurt samples and stirred cultured camel milk samples with 2.00% BSSP and 2.00% corn starch (standard) as a thickening agent having the highest values. Also, it was apparent that increasing BSSP concentration from 0.4 to 2.0 % significantly (p<0.05) decreased moisture content in stirred fermented camel milk products, conversely increasing the total solids. This could be attributed to the fact that BSSP contained hydrocolloids. Hydrocolloids are hydrophilic in nature and are effective in absorbing water by physically trapping free water and confining them in the increased network density and chemically by facilitating a link with other water molecules, thus increasing the gel-water binding capacity (Abdelmoneim *et al.*, 2016; Bahrami *et al.*, 2013).

Total solids content is a significant factor in the determination of the viscosity and rheological characteristics of yoghurt. Studies have indicated that higher total solid solids may result in improved viscosity, thus positively impacting the quality of yoghurt (Lussier *et al.*, 2023). Total solids are the most important technological property that determines the physical stability of yoghurt gel structure. Insufficient total solids fortification leads to wheying off and weak body (Mwizerwa *et al.*, 2017).

The average total solid content values reported by other researchers are as follows: 12.2% (Adel *et al.*, 2011), 16.65% (Galeboe *et al.*, 2018), 9.24% (Ibrahim & El Zubeir, 2016), 11.83% (Mustafa, 2015)

### 3.1.2 Ash Content

The ash content values varied from 0.23% to 1.08% for stirred camel milk yoghurt as shown in Table 1 and 0.23% to 0.953% for stirred cultured camel milk as shown in Table 2. Addition of BSSP as a thickening agent significantly ( $p < 0.05$ ) increased the ash content of both stirred camel milk yoghurt and stirred cultured camel milk. Stirred camel milk yoghurt sample containing 2.0% BSSP and stirred cultured camel milk sample containing 2.0% BSSP had the highest percentage of ash content compared to other samples indicating that 2.0% BSSP is rich in minerals compared to corn starch when used as a thickener at the same concentration. Ash content is of significant importance as it serves as an indicator of the mineral composition within the product, thus contributing to its nutritional value and potential health benefits. The average ash content values reported by other researchers are as follows: 0.71% (Adel *et al.*, 2011), 1.13% (Galeboe *et al.*, 2018), 0.84% (Mustafa, 2015), 0.99% (Seifu, 2023).

## 3.2 Effect of BSSP on the Storage of Stirred fermented camel milk Products

### 3.2.1 pH

During storage, different treatments of BSSP had significant ( $p < 0.05$ ) effects on both stirred camel milk yoghurt and stirred cultured camel milk pH. During storage, the pH of the negative control in stirred camel milk yoghurt decreased from 5.02 to 4.88. The pH of the stirred camel milk yoghurt sample containing 2.0% corn starch decreased from 4.8 to 4.72. Samples treated with different concentrations of BSSP also decreased; the pH of the yoghurt sample containing 0.4% BSSP decreased from 4.81 to 4.70; the pH of the stirred camel milk yoghurt sample containing 0.8% BSSP decreased from 4.79 to 4.66; the pH of the stirred camel milk yoghurt sample containing 1.2% BSSP decreased from 4.77 to 4.61; the pH of the stirred camel milk yoghurt sample containing 1.6% BSSP decreased from 4.62 to 4.43; the pH of the stirred camel milk yoghurt sample containing 2.0% BSSP decreased from 4.59 to 4.40 after being stored for 21 days (Figure 1). In the case of stirred cultured camel milk, the pH of negative control decreased from 4.78 to 4.69. The pH of the stirred cultured camel milk sample containing 2.0% corn starch decreased from 4.69 to 4.60. Samples treated with different concentrations of BSSP decreased; the pH of the stirred cultured camel milk sample containing 0.4% BSSP decreased from 4.68 to 4.59; the pH of the stirred cultured camel milk sample containing 0.8% BSSP decreased from 4.67 to 4.55; the pH of the stirred cultured camel milk sample containing 1.2% BSSP decreased from 4.65 to 4.53; the pH of the stirred cultured camel milk sample containing 1.6% BSSP decreased from 4.62 to 4.51; the pH of the stirred cultured camel milk sample containing 2.0% BSSP decreased from 4.6 to 4.49 after being stored for 21 days (Figure 2). The continuous decrease in pH values could be attributed to conversion of lactose to lactic acid by lactic acid bacteria during fermentation and the subsequent storage period (1-21 days) (Al-Zoreky & Al-Otaibi, 2015; Hassaan *et al.*, 2019; Khalifa & Ibrahim, 2015; Oselu *et al.*, 2022; Sadler & Murphy, 2010). The decrease in pH can also be attributed to the ability of BSSP to retain water since it contains fibre thereby diluting the concentration of lactic acid and other organic acids produced by the starter cultures (Alqahtani *et al.*, 2023). There are several factors that can also reduce the pH of fermented milk including total soluble solids, additives and storage temperature (Dabija, 2018).

Increasing the concentration of BSSP from (0.4% to 2.0%), reduced the pH values in both stirred camel milk yoghurt and stirred cultured camel milk samples as shown in Figure 1 and 2. There was a progressive decrease of pH values for all stirred camel milk yoghurt samples and stirred cultured camel milk samples but the pH decrease for stirred camel milk yoghurt samples

containing 2.0% BSSP and stirred cultured camel milk sample containing 2.0% BSSP was significantly higher ( $p < 0.05$ ) than the negative control samples in both stirred camel milk yoghurt and stirred cultured camel milk samples. This could be due to the higher total solids content in stirred camel milk yoghurt samples containing 2.0% BSSP and stirred cultured camel milk sample containing 2.0% BSSP that may have hindered the movement of the liquid phase and influenced the activities of starter cultures to metabolize sugars, producing lactic acid and organic acids as the main metabolic end products (Galeboe *et al.*, 2018; Oselu *et al.*, 2022; Singh *et al.*, 2017).

These findings exhibited comparable trends to those observed by other researchers including; (Khalifa & Ibrahim, 2015; Oselu *et al.*, 2022). The pH of stirred camel milk yoghurt decreased with increasing fermentation time from 0-6 hours and storage time from 1 to 21 days. The negative control yoghurt sample experienced a reduction in pH levels from 6.68 to 4.47 within a span of 6 hours, further declining to 4.4 after a period of 21 days. All stirred camel milk yoghurt samples treated with stabilizers decreased in their pH values after 21 days of storage (Oselu *et al.*, 2022). The pH values of stirred camel milk yoghurt samples varied based on the type and amount of stabilizer added. Stabilizers caused a decrease in pH compared to the control samples, with the highest acidity observed in the stabilizer-treated stirred camel milk yoghurts. The decline in pH over time during storage was attributed to continued fermentation by lactic acid bacteria and the acidity contributed by the added stabilizers. This decline was more pronounced in stirred camel milk yoghurt containing stabilizers compared to the control (Khalifa & Ibrahim, 2015).

### **3.2.2 Syneresis**

All stirred camel milk yoghurt samples and stirred cultured camel milk samples had a significant ( $p < 0.05$ ) decrease in their syneresis percentage in comparison to the negative control samples that did not contain any thickening agent in stirred camel milk yoghurt and stirred cultured camel milk samples, indicating that combining gelling agents and thickening agents reduced susceptibility to syneresis of both stirred camel milk yoghurt and stirred cultured camel milk samples. Combining both gelling and thickening agents improves the texture of fermented milk texture by promoting coagulum formation and because of the solubilization ability of gelatin, it increases the intermolecular interactions with the thickening agents, improving the viscosity and holding the gels together reducing syneresis (Muliro *et al.*, 2013).

During storage, the susceptibility to syneresis of the yoghurt control sample increased from 92.5% to 94.7%; stirred camel milk yoghurt sample containing 0.4% BSSP increased from 42.2% to 75.8%; stirred camel milk yoghurt sample containing 0.8% BSSP increased from 30.3% to 65.2%; stirred camel milk yoghurt sample containing 1.2% BSSP increased from 21.3% to 60%; stirred camel milk yoghurt sample containing 1.6% BSSP increased from 13.5% to 53.7%; stirred camel milk yoghurt sample containing 2.0% BSSP increased from 1.2% to 49%; standard stirred camel milk yoghurt sample containing 2.0% corn starch increased from 0 to 42.8% after being stored for 21 days. As for stirred cultured camel milk susceptibility to syneresis increased during storage, the control sample increased from 91.3% to 95.3%; the stirred cultured camel milk sample containing 0.4% BSSP increased from 39.8% to 73.8%; the stirred cultured camel milk sample containing 0.8% BSSP increased from 29% to 65.8%; the stirred cultured camel milk sample containing 1.2% BSSP increased from 21% to 60.3%; the stirred cultured camel milk sample containing 1.6% BSSP increased from 16% to 52.7%; the stirred cultured camel milk sample containing 2.0% BSSP increased from 2.5% to 48.7%; the standard stirred cultured camel milk sample containing 2.0%

corn starch increased from 0% to 43.2% after being stored for 21 days. Susceptibility to syneresis was significantly ( $p < 0.05$ ) reduced with increase in concentration of BSSP from 0.4% to 2.0%. This can be explained by the highly branched structure of BSSP, which could easily interact with gelatin and form bonds with the protein matrix in camel milk and ions in aqueous phase and water (Rojas-Torres *et al.*, 2021). These current results corroborate the findings of (Muliro *et al.*, 2013; Oselu *et al.*, 2022; Rojas-Torres *et al.*, 2021; Wong *et al.*, 2020) who reported that increasing the concentration of stabilizers, using both gelling and thickening agents reduced syneresis. The decrease in syneresis of samples containing BSSP may also be attributed to gummy sugars in fibres, which can trap water. These findings also align with the outcomes reported by Arabshashi-Delouee *et al.*, (2020) for yoghurt containing flaxseed press cake and Alqahtani *et al.*, (2023) for yoghurt containing date press cake.

In summary, stirred camel milk yoghurt sample containing 2.0% BSSP and 2.0% corn starch and 2.0% BSSP and 2.0% BSSP in stirred cultured camel milk sample displayed significantly ( $p < 0.05$ ) lower syneresis percentage over the storage period (1-21 days) because of the higher concentration of thickening agents as compared to all the other stirred camel milk yoghurt and stirred cultured camel milk formulations. The negative control samples in both stirred camel milk yoghurt and stirred cultured camel milk samples exhibited high syneresis percentage over the storage period (1-21 days). It was worth noting that there was a sharp increase in syneresis percentage on day 21 of storage in all the stirred camel milk yoghurt and stirred cultured camel milk samples as illustrated in (Figure 3 and 4). This could be attributed to reduction in total solids content over the storage period due to breakdown of macromolecules (Mwizerwa *et al.*, 2017).

### 3.2.3 Viscosity

The viscosity values of stirred camel milk yoghurt and stirred cultured camel milk samples increased from day 1 to day 7 of storage, followed by a decline on day 14 and a sharp drop on day 21 of storage. There was no significant difference between the standard and stirred camel milk yoghurt sample containing 2.0% BSSP and stirred cultured camel milk sample containing 2.0 % BSSP but the standard had the highest viscosity values compared to BSSP when used as a thickening agent at the same concentration. This could be attributed to the gel -forming characteristics of corn starch that prompted interactions between casein micelles in camel milk and corn starch (Galeboe *et al.*, 2018; Oselu *et al.*, 2022; Singh *et al.*, 2017) and the difference in structure between BSSP and corn starch polysaccharide. The standard exhibited better and increased intermolecular associations forming stronger junction zones with gelatin that facilitated better gel reformation contributing to an increase in viscosity and reduced susceptibility to syneresis as compared to 2.0% BSSP mixed with 0.4% gelatin. It was worth noting that increasing the concentration of BSSP from (0.4 to 2.0%) significantly ( $p < 0.05$ ) increased the viscosity of both stirred camel milk yoghurt and stirred cultured camel milk during storage. This is because low levels of BSSP did not cover and interact with all the casein particles in camel milk to create sufficient electrostatic and steric repulsions to stabilize the dispersion but higher concentrations of BSSP could sufficiently interact with the casein micelles hence the higher viscosity values. The addition of hydrocolloids from butternut squash seeds influenced the viscoelastic parameters of the yoghurt, with significant differences observed in  $G'$  and  $G''$  values among samples containing hydrocolloids from butternut squash seeds. The frequency sweep results revealed changes in the gel-like behaviour of the yoghurt samples, with hydrocolloids from butternut squash seeds

contributing to the formation of a strong gel structures and interacting with milk proteins (Rojas-Torres *et al.*, 2021).

During storage, the viscosity values significantly ( $p < 0.05$ ) decreased from the day of processing to the end of storage. In stirred camel milk yoghurt, the viscosity of the negative control sample containing 0.4 % gelatin solely decreased from 34 cP to 26 cP; the viscosity of the stirred camel milk yoghurt sample containing 0.4% BSSP decreased from 179 cP to 85 cP; the viscosity of the stirred camel milk yoghurt sample containing 0.8% BSSP decreased from 422 cP to 221 cP; the viscosity of the stirred camel milk yoghurt sample containing 1.2% BSSP decreased from 619 cP to 362 cP; the viscosity of the stirred camel milk yoghurt sample containing 1.6% BSSP decreased from 827 cP to 443 cP; the viscosity of the stirred camel milk yoghurt sample containing 2.0% BSSP decreased from 1076 cP to 676 cP; the viscosity of the standard sample containing 2.0% corn starch decreased from 1404 cP to 850 cP after being stored for 21 days (Figure 5). As for stirred cultured camel milk, the viscosity of the negative control sample containing 0.4% gelatin solely decreased from 50 cP to 32 cP; the viscosity of the stirred cultured camel milk sample containing 0.4% BSSP decreased from 147 cP to 87 cP; the viscosity of the stirred cultured camel milk sample containing 0.8% BSSP decreased from 431 cP to 249 cP; the viscosity of the stirred cultured camel milk sample containing 1.2% BSSP decreased from 602 cP to 365 cP; the viscosity of the stirred cultured camel milk sample containing 1.6% BSSP decreased from 784 cP to 438 cP; the viscosity of the stirred cultured camel milk sample containing 2.0% BSSP decreased from 957 cP to 577 cP; the viscosity of the standard sample containing 2.0% corn starch decreased from 1282 cP to 739 cP following 21 days of storage (Figure 6).

From the results of the present study there was significant ( $p < 0.05$ ) difference in stirred camel milk yoghurt and stirred cultured camel milk samples containing thickening agents and the negative control samples containing only a gelling agent indicating the important role played by thickening agents in improving and increasing the viscosity of stirred fermented camel milk products. Viscosity was low in gels formed with either gelatin or corn starch alone but combining both gelling and thickening agents improved the viscosity and reduced syneresis of stirred fermented camel milk products (Muliro *et al.*, 2013). These findings exhibited similar trends with those observed other researchers reporting that increasing concentrations of stabilizers, increased viscosity (Khalifa & Ibrahim, 2015; Muliro *et al.*, 2013; Oselu *et al.*, 2022; Rojas-Torres *et al.*, 2021; Wong *et al.*, 2020). BSSP also contains fibre and several studies have also indicated that fortifying yoghurt with various dietary fibres enhances the viscosity and thickening characteristics of fermented milk (Alqahtani *et al.*, 2023; Karaca *et al.*, 2019; Varnaité *et al.*, 2022).

Incorporating hydrocolloids sourced from butternut squash seeds as a stabilizer resulted in a decrease in apparent yield stress across all yoghurt samples, indicating intensified crosslinking of milk proteins in acidified milk, thereby impacting the yoghurt's structure under shear conditions. Research has showed that yoghurt samples exhibited non-Newtonian flow behaviour, whereas the apparent viscosity decreased as the shear rate increased, indicating shear-thinning tendency. The decrease in viscosity with increasing shear rate was attributed to the aligned flow of stiff polymer molecules, which reduced interactions between adjacent polymer molecules, which reduced interactions between adjacent polymer chains and disrupted the casein micelle network during shear application. Yoghurt samples containing hydrocolloids showed a two-step yield stress, indicating changes in the structural properties of the yoghurt matrix. The activation energy values of the yoghurt samples increased with the addition of hydrocolloids suggesting a higher

temperature dependence of viscosity in samples with higher hydrocolloid content (Rojas-Torres *et al.*, 2021).

## 4. CONCLUSION

This study demonstrates the promising potential of utilizing discarded butternut squash (*Cucurbita moschata*) seeds as a thickening agent in the development of stirred fermented camel milk products. The inclusion of BSSP in stirred cultured camel milk and yoghurt samples resulted in significant ( $p < 0.05$ ) reduction in syneresis values and an increase in viscosity values compared to the control. Moreover, higher concentrations of BSSP proved to be effective in further enhancing viscosity and reducing syneresis these products. The incorporation of BSSP emerges as a viable and sustainable alternative, not only contributing to waste reduction by utilizing butternut squash seeds but also offering tangible benefits in terms of texture improvement in stirred fermented camel milk products. These results present an opportunity for the food industry, particularly those involved in the production of stirred fermented camel milk products, to consider innovative approaches for enhancing product quality and consumer satisfaction.

## REFERENCES

- Abdelmoneim, A. H., Sherif, A. M., & Sameh, K. A. (2016a). *Rheological Properties of Yoghurt Manufactured by using Different Types of Hydrocolloids*.  
<https://www.semanticscholar.org/paper/Rheological-Properties-of-Yoghurt-Manufactured-by-Abdelmoneim-Sherif/c4107d13e727cb48c7780eac274ede6b5e935a30>
- Adel, E., Mohamed Ahmed, I., Babiker, E., & Yagoub, A. E. (2011). Physicochemical, Microbiological and Sensory Characteristics of Yoghurt Produced from Camel Milk During Storage. *Electronic Journal of Environmental, Agricultural and Food Chemistry*, *10*, 2305–2313.
- Alqahtani, N. K., Alnemr, T. M., Alsalem, A. K., Alotaibi, M. M., & Mohammed, M. (2023). Experimental Investigation and Modeling for the Influence of Adding Date Press Cake on Drinkable Yogurt Quality. *Foods*, *12*(6), 1219. <https://doi.org/10.3390/foods12061219>
- Al-Zoreky, N., & Al-Otaibi, M. (2015). Suitability of camel milk for making yogurt. *Food Science and Biotechnology*, *24*, 601–606. <https://doi.org/10.1007/s10068-015-0078-z>
- Bahrami, M., Ahmadi, D., Alizadeh, M., & Hosseini, F. (2013). Physicochemical and Sensorial Properties of Probiotic Yogurt as Affected by Additions of Different Types of Hydrocolloid. *Korean Journal for Food Science of Animal Resources*, *33*(3), 363–368. <https://doi.org/10.5851/kosfa.2013.33.3.363>
- Berhe, T., Seifu, E., Ipsen, R., Kurtu, M. Y., & Hansen, E. B. (2017). Processing Challenges and Opportunities of Camel Dairy Products. *International Journal of Food Science*, *2017*, 1–8. <https://doi.org/10.1155/2017/9061757>
- Cheng, H., & Hu, Y. (2010). Municipal solid waste (MSW) as a renewable source of energy: Current and future practices in China. *Bioresource Technology*, *101*(11), 3816–3824. <https://doi.org/10.1016/j.biortech.2010.01.040>
- Dabija, A. (2018, June 20). Physicochemical and sensory properties of yogurt with seabuckthorn powder, rosehip powder and grape seed extract during storage. 18th International Multidisciplinary Scientific GeoConference SGEM2018. <https://doi.org/10.5593/sgem2018/6.2/S25.057>
- De Kruif C.G., Tuinier R. Polysaccharide protein interactions. Food

- Hydrocolloids. 2001;15(4):555–563. doi: 10.1016/S0268-005X(01)00076-5
- Galeboe, O., Seifu, E., & Sekwati-Monang, B. (2018). *Production of Stirred camel milk yoghurt: Physicochemical and Microbiological Quality and Consumer Acceptability*. <https://doi.org/10.7455/ijfs/7.2.2018.a5>
- Hashim, I. B., Khalil, A. H., & Habib, H. (2009). Quality and acceptability of a set-type yogurt made from camel milk. *Journal of Dairy Science*, 92(3), 857–862. <https://doi.org/10.3168/jds.2008-1408>
- Hassaan, M., Ahmed, R., & A. Khider, M. (2019). Low Fat Flavored Stirred Yogurt Treated with different Stabilizers for Improving Its Properties and Quality. *Journal of Food and Dairy Sciences*, 10(12), 473–478. <https://doi.org/10.21608/jfds.2019.71364>
- Hematyar, N., Samarin, A. M., Poorazarang, H., Elhamirad, A., & Branch, S. (2012). *Effect of Gums on Yogurt Characteristics*. <https://www.semanticscholar.org/paper/Effect-of-Gums-on-Yogurt-Characteristics-Hematyar-Samarin/7493ed5da013c14741fba370cc048de10105301a>
- Ho, T. M., Zhao, J., & Bansal, N. (2022). Acid Gelation Properties of Camel Milk—Effect of <https://doi.org/10.1007/s11947-022-02890-5>
- Ibrahim, S. A., & El Zubeir, I. E. M. (2016). Processing, composition and sensory characteristic of yoghurt made from camel milk and camel–sheep milk mixtures. *Small Ruminant Research*, 136, 109–112. <https://doi.org/10.1016/j.smallrumres.2016.01.014>
- Jasim, M. A. J., Salih, G. M. S., Tikrit University, Hamk, M. L. H., & Slemani Polytechnic University. (2018). The Effect of Arabic Gum on Physicochemical and Sensory Properties of Camel Milk, Yogurt. *Journal of Zankoy Sulaimani - Part A, 2nd Int. Conf. AGR* (Special Issue), 97–104. <https://doi.org/10.17656/jzs.10656>
- Javidi F., Razavi S.M.A., Behrouzian F., Alghooneh A. The influence of basil seed gum, guar gum and their blend on the rheological, physical and sensory properties of low-fat ice cream. *Food Hydrocolloids*. 2016;52:625–633. doi: 10.1016/j.foodhyd.2015.08.006.
- Jiao, J., Li, Z.-G., Gai, Q.-Y., Li, X.-J., Wei, F.-Y., Fu, Y.-J., & Ma, W. (2014). Microwave-assisted aqueous enzymatic extraction of oil from pumpkin seeds and evaluation of its physicochemical properties, fatty acid compositions and antioxidant activities. *Food Chemistry*, 147, 17–24. <https://doi.org/10.1016/j.foodchem.2013.09.079>
- Karaca, O., Güzeler, N., Tangüler, H., Yaşar, K., & Akın, M. (2019). Effects of Apricot Fibre on the Physicochemical Characteristics, the Sensory Properties and Bacterial Viability of Nonfat Probiotic Yoghurts. *Foods*, 8(1), 33. <https://doi.org/10.3390/foods8010033>
- Kavas, N. (2016). Yoghurt production from camel (*Camelus dromedarius*) milk fortified with samphire molasses and different colloids. *Mljekarstvo / Dairy*, 66, 34–47. <https://doi.org/10.15567/mljekarstvo.2016.0104>
- Khalifa, S., & Ibrahim, A. (2015). Influence of addition modified starches as stabilizer on physicochemical and textural properties of camel’s milk yoghurt. *Zagazig Journal of Food and Dairy Research*.
- Li, H. (2020). Evaluation of bioactivity of butternut squash (*Cucurbita moschata* D.) seeds and skin. *Food Science & Nutrition*, 8(7), 3252–3261. <https://doi.org/10.1002/fsn3.1602>
- Liu, X., Gao, X., Wang, W., Zheng, L., Zhou, Y., & Sun, Y. (2012). Pilot-scale anaerobic co-digestion of municipal biomass waste: Focusing on biogas production and GHG reduction. *Renewable Energy*, 44, 463–468. <https://doi.org/10.1016/j.renene.2012.01.092>
- Lunardello, K. A., Yamashita, F., De Toledo Benassi, M., & De Rensis, C. M. V. B. (2012). The

- physicochemical characteristics of nonfat set yoghurt containing some hydrocolloids. *International Journal of Dairy Technology*, 65(2), 260–267.
- Lussier, N., Gilbert, A., St-Gelais, D., & Turgeon, S. L. (2023). Effect of the Heat Exchanger Type on Stirred Yogurt Properties Formulated at Different Total Solids and Fat Contents. *Dairy*, 4(1), Article 1. <https://doi.org/10.3390/dairy4010008>
- Muhammad Imran, H. A. S. (2014). Effects of Polymerized Whey Proteins Isolates on the Quality of Stirred Yoghurt made from Camel Milk. *Journal of Food Processing & Technology*, 05(07). <https://doi.org/10.4172/2157-7110.1000350>
- Muliro, P., Shalo, P., & Kutima, P. M. (2013). Optimization of camel milk coagulum formation and consumer preference. *Undefined*. <https://www.semanticscholar.org/paper/Optimization-of-camel-milk-coagulum-formation-and-Muliro-Shalo>. <http://dx.doi.org/10.14303/ajfst.2013.036>
- Mustafa, E. (2015). Comparison between the physiochemical attributes of yogurt processed from camel milk and that processed from cow milk and the effect of storage period on pH and acidity.
- Mwizerwa, H., Ooko Abong, G., Okoth, M., Ongol, M., Onyango, C., & Thavarajah, P. (2017). Effect of Resistant Cassava Starch on Quality Parameters and Sensory Attributes of Yoghurt. *Current Research in Nutrition and Food Science Journal*, 5(3), 353–367. <https://doi.org/10.12944/CRNFSJ.5.3.21>
- Oselu, S., Ebere, R., Huka, G., Musalia, L., Marete, E., Mathara, J. M., Mwobobia, F., & Arimi, J. M. (2022). Production and characterisation of stirred camel milk yoghurt containing different types of stabilising agents. *Heliyon*, 8(11), e11816. <https://doi.org/10.1016/j.heliyon.2022.e11816>
- Pirsa, S., & Hafezi, K. (2023). Hydrocolloids: Structure, preparation method, and application in food industry. *Food Chemistry*, 399, 133967. <https://doi.org/10.1016/j.foodchem.2022.133967>
- Rojas-Torres, S. A., Quintana, S. E., & García-Zapateiro, L. A. (2021). Natural Yogurt Stabilized with Hydrocolloids from Butternut Squash (*Cucurbita moschata*) Seeds: Effect on Physicochemical, Rheological Properties and Sensory Perception. *Fluids*, 6(7), 251. <https://doi.org/10.3390/fluids6070251>
- Sadler, G. D., & Murphy, P. A. (2010). pH and Titratable Acidity. In S. S. Nielsen (Ed.), *Food Analysis* (pp. 219–238). Springer US. [https://doi.org/10.1007/978-1-4419-1478-1\\_13](https://doi.org/10.1007/978-1-4419-1478-1_13)
- Seifu, E. (2023). Camel milk products: Innovations, limitations and opportunities. *Food Production, Processing and Nutrition*, 5(1), 15.
- Singh, R., Mal, G., Kumar, D., Patil, N. V., & Pathak, K. M. L. (2017). Camel Milk: An Important Natural Adjuvant. *Agricultural Research*, 6(4), 327–340. <https://doi.org/10.1007/s40003-017-0284-4>
- Szterk, A., Roszko, M., Sosińska, E., Derewiaka, D., & Lewicki, P. P. (2010). Chemical Composition and Oxidative Stability of Selected Plant Oils. *Journal of the American Oil Chemists' Society*, 87(6), 637–645. <https://doi.org/10.1007/s11746-009-1539-4>
- Varnaitė, L., Keršienė, M., Šipailienė, A., Kazernavičiūtė, R., Venskutonis, P. R., & Leskauskaitė, D. (2022). Fiber-Rich Cranberry Pomace as Food Ingredient with Functional Activity for Yogurt Production. *Foods*, 11(5), 758. <https://doi.org/10.3390/foods11050758>
- Wang, L., Liu, F., Wang, A., Yu, Z., Xu, Y., & Yang, Y. (2017). Purification, characterization and bioactivity determination of a novel polysaccharide from pumpkin (*Cucurbita moschata*) seeds. *Food Hydrocolloids*, 66, 357–364. <https://doi.org/10.1016/j.foodhyd.2016.12.003>

- Wong, S.-S., Wicklund, R., Bridges, J., Whaley, J., & Koh, Y. B. (2020). Starch swelling behavior and texture development in stirred yogurt. *Food Hydrocolloids*, 98, 105274.
- Wusigale, Liang L., Luo Y. Casein and pectin: Structures, interactions, and applications. *Trends in Food Science & Technology*. 2020; 97:391–403. doi: 10.1016/j.tifs.2020.01.027